

|                    |  |                  |  |
|--------------------|--|------------------|--|
| <b>Client Name</b> |  | <b>Client ID</b> |  |
|--------------------|--|------------------|--|

|                       |  |               |  |
|-----------------------|--|---------------|--|
| <b>Effective Date</b> |  | <b>Author</b> |  |
|-----------------------|--|---------------|--|

|                           |  |
|---------------------------|--|
| <b>Type of Assessment</b> |  |
|---------------------------|--|

**Dimension 1: Acute Intoxication and/or Withdrawal Potential**

- No withdrawal risk (Level 0.5)
- Physiologically dependent on opiates and requires Opioid Maintenance Therapy (OTP Level 1)
- Withdrawal, if present, is manageable at Level 1-WM (Level 1)
- Withdrawal, if present, is manageable at Level 2-WM (Level 2.1)
- Withdrawal, if present, is manageable at Level 2-WM (Level 2.5)
- Withdrawal, if present, is currently receiving Level 1-WM or 3.2-WM services (Level 3.1)
- Withdrawal, if present, is manageable at Level 3.2-WM (Level 3.3)
- Withdrawal, if present, is manageable at Level 3.2-WM (Level 3.5)
- Withdrawal, if present, is manageable at Level 3.7-WM (Level 3.7)
- Withdrawal requires Level 4-WM (Level 4)

**General**

|              |  |
|--------------|--|
| <b>Level</b> |  |
|--------------|--|

|                        |  |
|------------------------|--|
| <b>Documented Risk</b> |  |
|------------------------|--|

Comments

|  |
|--|
|  |
|--|

**Dimension 2: Biomedical Conditions and Complications**

- None or very stable (Level 0.5)
- None or manageable with outpatient medical monitoring (OTP Level 1)
- None or very stable, or the patient receiving concurrent medical monitoring (Level 1)
- None or not a distraction from treatment. Such problems are manageable at Level 2.1 (Level 2.1)
- None or not sufficient to distract from treatment. Such problems are manageable at Level 2.5 (Level 2.5)
- None or stable, or the patient is receiving concurrent medical monitoring (Level 3.1)
- None or stable, or receiving concurrent medical treatment (Level 3.3)
- None or stable, or receiving concurrent medical treatment (Level 3.5)
- None or stable, or receiving concurrent medical treatment (Level 3.7)
- Requires 24-hour medical and nursing care in a hospital (Level 4)

**General**

|              |  |
|--------------|--|
| <b>Level</b> |  |
|--------------|--|

|                        |  |
|------------------------|--|
| <b>Documented Risk</b> |  |
|------------------------|--|

Comments

|  |
|--|
|  |
|--|

**Dimension 3: Emotional, Behavioral or Cognitive Conditions and Complications**

- None or very stable (Level 0.5)
- None or manageable with outpatient structured monitoring (OPT Level 1)
- None or very stable, or the patient receiving concurrent mental health monitoring (Level 1)
- Mild severity, with the potential to distract from recovery; needs monitoring (Level 2.1)
- Mild to moderate severity, with potential to distract from recovery; needs stabilization (Level 2.5)
- None or minimal, not distracting from recovery (Level 3.1)
- Mild to moderate severity; needs structure to focus on recovery (Level 3.3)
- Inability to control impulses or unstable and dangerous signs/symptoms require stabilization and a 23-hour setting (Level 3.5)
- Moderate Severity; needs a 24-hour structured setting (Level 3.7)
- Requires 24-hour psychiatric care and concomitant addiction treatment (Level 4)

**General**

|              |  |
|--------------|--|
| <b>Level</b> |  |
|--------------|--|

|                        |  |
|------------------------|--|
| <b>Documented Risk</b> |  |
|------------------------|--|

Comments

|  |
|--|
|  |
|--|

**Dimension 4: Readiness to Change**

- Willing to explore how current alcohol, tobacco, other drug or medication and/or high risk behaviors may affect personal goals (Level 0.5)
- Ready to change the negative effects of opioid use, but not ready for total abstinence from illicit prescription or non-prescription drug use (OTP Level 1)
- Ready for recovery but needs motivation and monitoring strategies to strength readiness; or needs ongoing monitoring and disease management; or high severity in this dimensions but no in other dimensions (Level 1)
- Has variable engagement in treatment, ambivalence, or lack of awareness of the substance use or mental health problem, and requires a structured program several times a week (Level 2.1)
- Has poor engagement in treatment, significant ambivalence, or a lack of awareness of the substance use or mental health problem, requiring a near-daily structured program or intensive engagement services (Level 2.5)
- Open to recovery, but needs a structured environment to maintain therapeutic gains (Level 3.1)
- Has little awareness and needs interventions available only in Level 3.3 to stay in treatment (Level 3.3)
- Has marked difficulty with, or opposition to, treatment, with dangerous consequences (Level 3.5)
- Low interest in treatment and impulse control is poor, despite negative consequences; needs motivating strategies available in a 24 hour structured setting (Level 3.7)
- Requires 24-hour psychiatric care and concomitant addiction treatment (Level 4)

**General**

|              |  |
|--------------|--|
| <b>Level</b> |  |
|--------------|--|

|                        |  |
|------------------------|--|
| <b>Documented Risk</b> |  |
|------------------------|--|

Comments

|  |
|--|
|  |
|--|

**Dimension 5: Relapse, Continued Use, Continued Problem Potential**

- Needs an understanding of, or skills to change, current alcohol, tobacco, other drug, or medication use patterns, and/or high risk behavior (Level 0.5)
- At high risk of relapse or continued use without OTP and structured therapy (OPT Level 1)
- Able to maintain abstinence or control use an/or addictive behaviors and pursue recovery or motivational foals with minimal support (Level 1)
- Intensification of addiction or mental health symptoms indicate a high likelihood of relapse or continued use or continued problems without close monitoring and support several times a week (Level 2.1)
- Intensification of addiction or mental health systems, despite active participation in level 1 or level 2.1 program, indicates a high likelihood of relapse or continued use or continued problems without near-daily monitoring and support (Level 2.5)
- Understands relapse but needs structure to maintain therapeutic gains (Level 3.1)
- Has little awareness and needs interventions available only at Level 3.3 to prevent continued use, with imminent dangerous consequences, because of cognitive deficits or comparable dysfunction (Level 3.3)
- Has no recognition of the skills needed to prevent continued use, with imminently dangerous consequences (Level 3.5)
- Unable to control use, with imminent dangerous consequences despite active participation at less intensive levels of care (Level 3.7)
- Problems in this dimension do not qualify this person for Level 4 services (Level 4)

**General**

|              |  |
|--------------|--|
| <b>Level</b> |  |
|--------------|--|

|                        |  |
|------------------------|--|
| <b>Documented Risk</b> |  |
|------------------------|--|

Comments

|  |
|--|
|  |
|--|

**Dimension 6: Recovery/Living Environment**

- Social support system or significant others increase the risk of personal conflict about alcohol, tobacco, or other drug use (Level 0.5)
- Recovery environment is supportive and/or the person has skills to cope (OPT Level 1)
- Recovery environment is supportive and/or the person has skills to cope (Level 1)
- Recovery environment is not supportive, but with structure and support, the person can cope (Level 2.1)
- Recovery environment is not supportive, but with structure and support and relief from the home environment, the person can cope (Level 2.5)
- Environment is dangerous, but recovery is achievable if level 3.1 / 24 hour structure is available (Level 3.1)
- Environment is dangerous and person needs 24-hour structure to learn to cope (Level 3.3)
- Environment is dangerous and the person lack skills to cope outside of a highly structured 24 setting (Level 3.5)
- Environment is dangerous and the person lack skills to cope outside of a highly structured 24 setting (Level 3.7)
- Problems in this dimension do not qualify this person for Level 4 services (Level 4)

**General**

|              |  |
|--------------|--|
| <b>Level</b> |  |
|--------------|--|

|                        |  |
|------------------------|--|
| <b>Documented Risk</b> |  |
|------------------------|--|

Comments

|  |
|--|
|  |
|--|

**Dimension 6: Recovery/Living Environment**

|                    |                    |
|--------------------|--------------------|
| <b>Dimension 1</b> | <b>Dimension 2</b> |
| Level:             | Level:             |
| Risk:              | Risk:              |
|                    |                    |
| <b>Dimension 3</b> | <b>Dimension 4</b> |
| Level:             | Level:             |
| Risk:              | Risk:              |
|                    |                    |
| <b>Dimension 5</b> | <b>Dimension 6</b> |
| Level:             | Level:             |
| Risk:              | Risk:              |
|                    |                    |

**Final Placement Determination**

|                            |  |
|----------------------------|--|
| Indicated Level of Care/WM |  |
| Actual Level of Care/WM    |  |

**Comments**

**San Diego ASAM Final Determination**

**Final Placement Determination**

|                                       |  |
|---------------------------------------|--|
| Additional Indicated Level of Care/WM |  |
|---------------------------------------|--|

|                                   |  |
|-----------------------------------|--|
| Second Indicated Level of Care/WM |  |
|-----------------------------------|--|

|   |  |
|---|--|
| Additional Actual Level of Care/WM Decision |  |
|---|--|

If Actual LOC was not among those indicated, what is the reason for the difference?

|  |
|--|
|  |
|--|

If reason was "Other", explain:

|  |
|--|
|  |
|--|

If referral is being made but admission is expected to be delayed, what is the reason for delay?

|  |
|--|
|  |
|--|

If reason was "Other", explain:

|  |
|--|
|  |
|--|

**Immediate Need Profile Determination**

Outcome of Immediate Needs Profile

|  |
|--|
|  |
|--|

|                            |  |             |  |
|----------------------------|--|-------------|--|
| <b>Counselor Signature</b> |  | <b>Date</b> |  |
|----------------------------|--|-------------|--|

|   |  |
|---|--|
| <b>Counselor Printed Name &amp; Credentials</b> |  |
|---|--|

|                       |  |             |  |
|-----------------------|--|-------------|--|
| <b>LPHA Signature</b> |  | <b>Date</b> |  |
|-----------------------|--|-------------|--|

|  |  |
|--|--|
| <b>LPHA Printed Name &amp; Credentials</b> |  |
|--|--|